

HIV and Its Transmission FactSheet

HS02-034B (9-04)

Research has revealed a great deal of valuable medical, scientific, and public health information about the human immunodeficiency virus (HIV) and acquired immunodeficiency syndrome (AIDS). The ways in which HIV can be transmitted have been clearly identified. Unfortunately, false information or statements that are not supported by scientific findings continue to be shared widely through the Internet and popular press. Therefore, the Centers for Disease Control and Prevention (CDC) prepared this fact sheet to correct a few misperceptions about HIV.

How HIV is Transmitted

HIV is spread by sexual contact with an infected person, by sharing needles and/or syringes (primarily for drug injection) with someone who is infected, or, less commonly (and now very rarely in countries where blood is screened for HIV antibodies), through transfusions of infected blood or blood clotting factors. Babies born to HIV-infected women may become infected before or during birth or through breast-feeding after birth.

In the health care setting, workers have been infected with HIV after being stuck with needles containing HIV-infected blood or, less frequently, after infected blood gets into a worker's open cut or a mucous membrane (for example, the eyes or inside of the nose). There has been only one instance of patients being infected by a health care worker in the United States; this involved HIV transmission from one infected dentist to six patients. Investigations have been completed involving more than 22,000 patients of 63 HIV-infected physicians, surgeons, and dentists, and no other cases of this type of transmission have been identified in the United States.

Some people fear that HIV might be transmitted in other ways; however, no scientific evidence to support any of these fears has been found. If HIV were being transmitted through other routes (such as through air, water, or insects), the pattern of reported AIDS cases would be much different from what has been observed. For example, if mosquitoes could transmit HIV infection, many more young children and preadolescents would have been diagnosed with AIDS.

All reported cases suggesting new or potentially unknown routes of transmission are thoroughly investigated by state and local health departments with the assistance,

guidance, and laboratory support from CDC. No additional routes of transmission have been recorded, despite a national sentinel system designed to detect just such an occurrence.

The following paragraphs specifically address some of the common misperceptions about HIV transmission.

HIV in the Environment

Scientists and medical authorities agree that HIV does not survive well in the environment, making the possibility of environmental transmission remote. HIV is found in varying concentrations or amounts in blood, semen, vaginal fluid, breast milk, saliva, and tears. (See *Saliva, Tears, and Sweat*.) To obtain data on the survival of HIV, laboratory studies have required the use of artificially high concentrations of laboratory-grown virus. Although these unnatural concentrations of HIV can be kept alive for days or even weeks under precisely controlled and limited laboratory conditions, CDC studies have shown that drying of even these high concentrations of HIV reduces the amount of infectious virus by 90 to 99 percent within several hours. Since the HIV concentrations used in laboratory studies are much higher than those actually found in blood or other specimens, drying of HIV-infected human blood or other body fluids reduces the risk of environmental transmission to that which has been observed — essentially zero. Incorrect interpretations of conclusions drawn from laboratory studies have unnecessarily alarmed some people.

Results from laboratory studies should not be used to assess specific personal risk of infection because (1) the amount of virus studied is not found in human specimens or elsewhere in nature, and (2) no one has been identified as infected with HIV due to contact with an environmental surface. Additionally, HIV is unable to reproduce outside its living host (unlike many bacteria or fungi, which may do so under suitable conditions), except under laboratory conditions; therefore, it does not spread or maintain infectiousness outside its host.



Businesses and Other Settings

There is no known risk of HIV transmission to co-workers, clients, or consumers from contact in industries such as food-service establishments (see *HIV in the Environment*). Food-service workers, known to be infected with HIV, need not be restricted from work unless they have other infections or illnesses (such as diarrhea or hepatitis A) for which any food-service worker, regardless of HIV infection status, should be restricted. CDC recommends that all food-service workers follow recommended standards and practices of good personal hygiene and food sanitation.

In 1985, CDC issued routine precautions that all personal-service workers (such as hairdressers, barbers, cosmetologists, and massage therapists) should follow, even though there is no evidence of transmission from a personal-service worker to a client or vice versa. Instruments that are intended to penetrate the skin (such as tattooing and acupuncture needles, ear piercing devices) should be used once and disposed of or thoroughly cleaned and sterilized. Instruments not intended to penetrate the skin but which may become contaminated with blood (for example, razors) should be used for only one client and disposed of or thoroughly cleaned and disinfected after each use. Personal-service workers can use the same cleaning procedures that are recommended for health care institutions.

CDC knows of no instances of HIV transmission through tattooing or body piercing, although hepatitis B virus has been transmitted during some of these practices. One case of HIV transmission from acupuncture has been documented. Body piercing (other than ear piercing) is relatively new in the United States, and the medical complications for body piercing appear to be greater than for tattoos. Healing of piercing generally will take weeks, and sometimes even months, and the pierced tissue could conceivably be abraded (torn or cut) or inflamed even after healing. Therefore, a theoretical HIV transmission risk does exist if the unhealed or abraded tissues come into contact with an infected person's blood or other infectious body fluid. Additionally, HIV could be transmitted if instruments contaminated with blood are not sterilized or disinfected between clients.

Households

Although HIV has been transmitted between family members in a household setting, this type of transmission is very rare. These transmissions are believed to have resulted from contact between skin or mucous membranes and infected blood. To prevent even such rare occurrences, precautions, as described in previously published guidelines, should be taken in all settings “including the

home”, to prevent exposures to the blood of persons who are HIV infected, at risk for HIV infection, or whose infection and risk status are unknown. For example,

- Gloves should be worn during contact with blood or other body fluids that could possibly contain visible blood, such as urine, feces, or vomit.
- Cuts, sores, or breaks on both the caregiver's and patient's exposed skin should be covered with bandages.
- Hands and other parts of the body should be washed immediately after contact with blood or other body fluids, and surfaces soiled with blood should be disinfected appropriately.
- Practices that increase the likelihood of blood contact, such as sharing of razors and toothbrushes, should be avoided.
- Needles and other sharp instruments should be used only when medically necessary and handled according to recommendations for health-care settings. (Do not put caps back on needles by hand or remove needles from syringes.) Dispose of needles in puncture-proof containers.

Saliva, Tears, and Sweat

HIV has been found in saliva and tears in very low quantities from some AIDS patients. It is important to understand that finding a small amount of HIV in a body fluid does not necessarily mean that HIV can be transmitted by that body fluid. HIV has not been recovered from the sweat of HIV-infected persons. Contact with saliva, tears, or sweat has never been shown to result in transmission of HIV.

Insects

From the onset of the HIV epidemic, there has been concern about transmission of the virus by biting and bloodsucking insects. However, studies conducted by researchers at CDC and elsewhere have shown no evidence of HIV transmission through insects — even in areas where there are many cases of AIDS and large populations of insects such as mosquitoes. Lack of such outbreaks, despite intense efforts to detect them, supports the conclusion that insects do not transmit HIV.

The results of experiments and observations of insect biting behavior indicate that when an insect bites a person, it does not inject its own or a previously bitten person's or animal's blood into the next person bitten. Rather, it injects saliva, which acts as a lubricant or anticoagulant so the insect can feed efficiently. Such diseases as yellow fever

and malaria are transmitted through the saliva of specific species of mosquitoes. However, HIV lives for only a short time inside an insect and, unlike organisms that are transmitted via insect bites, HIV does not reproduce (and does not survive) in insects. Thus, even if the virus enters a mosquito or another sucking or biting insect, the insect does not become infected and cannot transmit HIV to the next human it feeds on or bites. HIV is not found in insect feces.

There is also no reason to fear that a biting or bloodsucking insect, such as a mosquito, could transmit HIV from one person to another through HIV-infected blood left on its proboscis. Two factors serve to explain why this is so — first, infected people do not have constant, high levels of HIV in their bloodstreams and, second, insect parts do not retain large amounts of blood on their surfaces. Further, scientists who study insects have determined that biting insects normally do not travel from one person to the

next immediately after ingesting blood. Rather, they fly to a resting place to digest this blood meal.

For more information...

CDC National AIDS Hotline:

1-800-342-AIDS

Spanish: 1-800-344-SIDA

Deaf: 1-800-243-7889

Internet Resources:

<http://www.cdc.gov/nchstp/od/nchstp.html>

<http://www.cdc.gov/hiv>

<http://www.cdcnpin.org>

Disclaimer: Information contained in this training program is considered accurate at time of publication.